arXiv:1109.1658v1 [math.LO] 8 Sep 2011

ORTHOCOMPLEMENTED WEAK TENSOR PRODUCTS

BORIS ISCHI

Abstract. Let $\mathcal{L}_1$ and $\mathcal{L}_2$ be complete atomistic lattices. In a previous paper, we have defined a set $\mathsf{S} = \mathsf{S}(\mathcal{L}_1, \mathcal{L}_2)$ of complete atomistic lattices, the elements of which are called weak tensor products of $\mathcal{L}_1$ and $\mathcal{L}_2$. S is defined by means of three axioms, natural regarding the description of some compound systems in quantum logic. It has been proved that S is a complete lattice. The top element of S, denoted by $\mathcal{L}_1 \varovee \mathcal{L}_2$, is the tensor product of Fraser whereas the bottom element, denoted by $\mathcal{L}_1 \varowedge \mathcal{L}_2$, is the *box product* of Grätzer and Wehrung. With some additional hypotheses on $\mathcal{L}_1$ and $\mathcal{L}_2$ (true for instance if $\mathcal{L}_1$ and $\mathcal{L}_2$ are moreover orthomodular with the covering property) we prove that S is a singleton if and only if $\mathcal{L}_1$ or $\mathcal{L}_2$ is distributive, if and only if $\mathcal{L}_1 \varovee \mathcal{L}_2$ has the covering property. Our main result reads: $\mathcal{L} \in \mathsf{S}$ admits an orthocomplementation if and only if $\mathcal{L} = \mathcal{L}_1 \varowedge \mathcal{L}_2$. At the end, we construct an example $\mathcal{L}_1 \mathbin{\bigcirc\!\!\!\!\!\Downarrow} \mathcal{L}_2$ in S which has the covering property.

1. Introduction

Let Σ be a non-empty set. By a *simple closure space* $\mathcal{L}$ on Σ, we mean a set of subsets of Σ, ordered by set-inclusion, closed under arbitrary set-intersections (*i.e.*, for all $\omega \subseteq \mathcal{L}$, $\bigcap \omega \in \mathcal{L}$), and containing Σ, $\emptyset$, and all singletons. We denote the bottom ($\emptyset$) and top (Σ) elements by 0 and 1 respectively. For $p \in \Sigma$, we identify p with $\{p\} \in \mathcal{L}$. Hence $p \bigcup q$ stands for $\{p, q\}$.

Let $\mathcal{L}$ be a simple closure space on a (nonempty) set Σ. Then $\mathcal{L}$ is a complete atomistic lattice, the atoms of which correspond to the points (*i.e.*, singletons) of Σ. Note that if $A \subseteq \Sigma$, then $\bigvee_{\mathcal{L}}(A) = \bigcap\{b \in \mathcal{L} \mid A \subseteq b\}$. Conversely, let $\mathcal{L}$ be a complete atomistic lattice. Let Σ denote the set of atoms of $\mathcal{L}$, and, for each $a \in \mathcal{L}$, let $\Sigma[a]$ denote the set of atoms under a. Then $\{\Sigma[a] \mid a \in \mathcal{L}\}$ is a simple closure space on Σ, isomorphic to $\mathcal{L}$. For simplicity, we shall from now on deal only with simple closure spaces instead of complete atomistic lattices.

Let $\mathcal{L}_1$, $\mathcal{L}_2$ and $\mathcal{L}$ be simple closure spaces on Σ_1, Σ_2 and Σ respectively. Then, $\mathcal{L}$ is a *weak tensor product* of $\mathcal{L}_1$ and $\mathcal{L}_2$ if

(P1) $\Sigma = \Sigma_1 \times \Sigma_2$
(P2) $a_1 \times \Sigma_2 \cup \Sigma_1 \times a_2 \in \mathcal{L}$, $\forall\ a_i \in \mathcal{L}_i$
(P3) for all $p_1 \in \Sigma_1$ and $A_2 \subseteq \Sigma_2$, $p_1 \times A_2 \in \mathcal{L}$ implies $A_2 \in \mathcal{L}_2$
for all $p_2 \in \Sigma_2$ and $A_1 \subseteq \Sigma_1$, $A_1 \times p_2 \in \mathcal{L}$ implies $A_1 \in \mathcal{L}_1$

Let $\mathsf{S}(\mathcal{L}_1, \mathcal{L}_2)$ denote the set of weak tensor products of $\mathcal{L}_1$ and $\mathcal{L}_2$, ordered by set-inclusion. Write $\pi_i : \Sigma_1 \times \Sigma_2 \to \Sigma_i$ the projection map defined by $\pi_i(p_1, p_2) = p_i$. Then, it is easy to check that

$$\mathcal{L}_1 \varowedge \mathcal{L}_2 = \left\{ \bigcap \omega \mid \omega \subseteq \{a_1 \times \Sigma_2 \cup \Sigma_1 \times a_2 \mid a_i \in \mathcal{L}_i\},\ \omega \neq \emptyset \right\}$$

2010 *Mathematics Subject Classification.* Primary 06C15; Secondary 81P10, 06B23.
Key words and phrases. Complete atomistic ortholattice, tensor product, quantum logic.

is the bottom element of $\mathsf{S}(\mathcal{L}_1, \mathcal{L}_2)$ whereas

$$\mathcal{L}_1 \varovee \mathcal{L}_2 = \Big\{ A \subseteq \Sigma_1 \times \Sigma_2 \mid \ \pi_2\left(p_1 \times \Sigma_2 \cap A\right) \in \mathcal{L}_2 \ \text{ and } \\ \pi_1\left(\Sigma_1 \times p_2 \cap A\right) \in \mathcal{L}_1, \ \forall\, p_i \in \Sigma_i \Big\}$$

is the top element of $\mathsf{S}(\mathcal{L}_1, \mathcal{L}_2)$ [5].

A cornerstone in quantum logic is the following theorem, by which the Hilbert space structure of quantum mechanics can be recovered from a certain number of axioms on the lattice $\mathcal{L}_S$ of experimental propositions concerning a physical system S [7] (see [6], Theorems 34.5 and 34.9). Let E be a vector space over a $^*-$division ring $\mathbb{K}$. A *Hermitian form* f is a mapping f of $E \times E$ onto $\mathbb{K}$, linear in the left variable (*i.e.*, $f(\lambda_1 x_1 + \lambda_2 x_2, y) = \lambda_1 f(x_1, y) + \lambda_2 f(x_2, y)$, where $x_i, y \in E$ and $\lambda_i \in \mathbb{K}$), such that $f(x,y)^* = f(y,x)$ and $f(x,x) = 0$ implies $x = 0$. Let V be a subspace of E. Put $V^\perp = \{y \in E \mid f(x,y) = 0 \ \forall x \in V\}$ and say that V is $E-$*closed* if $V = V^{\perp\perp}$.

Theorem 1.1. *If $\mathcal{L}_S$ is an irreducible orthocomplemented simple closure space with the covering property, and of length ≥ 4, then there is a $^*-$division ring $\mathbb{K}$ and a vector space E over $\mathbb{K}$ with an Hermitian form such that $\mathcal{L}_S$ is ortho-isomorphic to the lattice $\mathsf{P}(E)$ of $E-$closed subspaces of E. Moreover, if $\mathbb{K} = \mathbb{R}$ or $\mathbb{C}$ with the usual involution, then E is a Hilbert space if and only if $\mathcal{L}$ is moreover orthomodular.*

For some compound quantum systems, called *separated systems*, it can be shown that $\mathcal{L}_S$, which we henceforth denote by $\mathcal{L}_{S_{sep}}$, must satisfy the three axioms P1-P3 defining weak tensor products (see [5] and references herein) and cannot be isomorphic to the lattice of closed subspaces of a Hilbert space [2]. As consequence, some hypotheses of Theorem 1.1 fail in $\mathcal{L}_{S_{sep}}$.

In [1], Aerts proposed $\mathcal{L}_1 \varowedge \mathcal{L}_2$ as a model for $\mathcal{L}_{S_{sep}}$. For $\mathcal{L}_1$ and $\mathcal{L}_2$ orthocomplemented simple closure spaces (in which case, $\mathcal{L}_1 \varowedge \mathcal{L}_2$ is an orthocomplemented simple closure space), Aerts proved that if $\mathcal{L}_1 \varowedge \mathcal{L}_2$ has the covering property or is orthomodular, then $\mathcal{L}_1$ or $\mathcal{L}_2$ is a power set [1]. As a consequence, according to Aerts, the covering property and orthomodularity do not hold in $\mathcal{L}_{S_{sep}}$. A similar conclusion concerning orthomodularity was obtained by Pulmannová in [9].

Here we argue that no reasonable model for $\mathcal{L}_{S_{sep}}$ admits an orthocomplementation. On the other hand, in case $\mathcal{L}_1$ and $\mathcal{L}_2$ satisfy the axioms of theorem 1.1, we provide a natural model for $\mathcal{L}_{S_{sep}}$ which has the covering property. We proceed as follows: following [8] and [1], we assume that $\mathcal{L}_1$, $\mathcal{L}_2$ and $\mathcal{L}_{S_{sep}}$ are simple closure spaces. Let $\mathcal{L} \in \mathsf{S}(\mathcal{L}_1, \mathcal{L}_2)$. We prove that if $\mathcal{L}_1$ and $\mathcal{L}_2$ are orthocomplemented simple closure spaces with the covering property, then $\mathcal{L}$ admits and orthocomplementation if and only if $\mathcal{L} = \mathcal{L}_1 \varowedge \mathcal{L}_2$. We conclude by a simple physical argument given in [4] which shows that certainly $\mathcal{L}_1 \varowedge \mathcal{L}_2 \subsetneqq \mathcal{L}_{S_{sep}}$.

The rest of the paper is organized as follows. In Section 2, we fix some basic terminology and notation. We define a set $\mathsf{S} \equiv \mathsf{S}(\mathcal{L}_1, \cdots, \mathcal{L}_n)$ of n-fold weak tensor products.

Further, with some additional hypotheses on each $\mathcal{L}_i$ (true for instance if $\mathcal{L}_i$ is moreover orthocomplemented with the covering property) we prove that S is a singleton (*i.e.*, $\varowedge_i \mathcal{L}_i = \varovee_i \mathcal{L}_i$) if and only if at most one $\mathcal{L}_i$ is not a power set (Section 3) if and only if $\varowedge_i \mathcal{L}_i$ or $\varovee_i \mathcal{L}_i$ has the covering property (Section 5). Note that for the last condition concerning $\varovee_i \mathcal{L}_i$, we need to make an additional assumption on each $\mathcal{L}_i$.

Finally, Section 4 is devoted to our main result and Section 6 to the example mentioned above.

2. Main definitions

In this section we give our main definitions. We start with some background material and basic notations used in the sequel. Parts of this section are taken directly from [5].

Definition 2.1.

- A lattice $\mathcal{L}$ with 0 and 1 is *orthocomplemented* if there is a unary operation $^\perp$ (orthocomplementation), also denoted by $'$, such that for all a, $b \in \mathcal{L}$, $(a^\perp)^\perp = a$, $a \leq b$ implies $b^\perp \leq a^\perp$, and $a \vee a^\perp = 1$.
- An orthocomplemented lattice is *orthomodular* if for all a, $b \in \mathcal{L}$, $a \leq b$ implies $b = a \vee (b \wedge a^\perp)$.
- A lattice with 0 has the *covering property* if for any atom p and any $a \in \mathcal{L}$, $p \wedge a = 0$ implies that $p \vee a$ covers a (in symbols $p \vee a \gtrdot a$).
- A lattice $\mathcal{L}$ with 0 and 1 is called a *DAC-lattice* if $\mathcal{L}$ and its dual $\mathcal{L}^*$ (defined by the converse order-relation) are atomistic with the covering property [6]. We say that a lattice $\mathcal{L}$ with 1 is *coatomistic* if the dual $\mathcal{L}^*$ is atomistic.
- 2 denotes the simple closure space isomorphic to the two-element lattice.
- Let $\mathcal{L}$ and $\mathcal{L}_1$ be simple closure spaces on Σ and Σ_1 respectively. We write $\mathsf{Aut}(\mathcal{L})$ for the group of automorphisms of $\mathcal{L}$. Note that any map $u \colon \mathcal{L} \to \mathcal{L}_1$ sending atoms to atoms induces a mapping from Σ to Σ_1, which we also call u. Thus, if $u \in \mathsf{Aut}(\mathcal{L})$, then for all $a \in \mathcal{L}$, $u(a) = \{u(p) \mid p \in a\}$.
- If $\mathcal{L}$ is orthocomplemented, for p, $q \in \Sigma$, we write $p \perp q$ if $p \in q^\perp$, where $q^\perp$ stands for $\{q\}^\perp$.
- If $\mathcal{H}$ is a complex Hilbert space, then $\Sigma_{\mathcal{H}}$ denotes the set of 1-dimensional subspaces of $\mathcal{H}$ and $\mathsf{P}(\mathcal{H})$ stands for the simple closure space on $\Sigma_{\mathcal{H}}$ isomorphic to the lattice of closed subspaces of $\mathcal{H}$. Moreover, we write $\mathsf{U}(\mathcal{H})$ for the set of automorphisms of $\mathsf{P}(\mathcal{H})$ induced by unitary maps on $\mathcal{H}$.
- Finally, we say that $\mathcal{L}$ (respectively $T \subseteq \mathsf{Aut}(\mathcal{L})$) is *transitive* if the action of $\mathsf{Aut}(\mathcal{L})$ (respectively the action of T) on Σ is transitive.

Remark 2.2. Note that an orthocomplemented atomistic lattice with the covering property is a DAC-lattice. Note also that in Theorem 1.1, if instead of orthocomplemented, the simple closure space is a DAC-lattice, then there is a pair of dual vector spaces such that a representation theorem similar to Theorem 1.1 holds (see [6], Theorem 33.7).

Definition 2.3. Let $\{\Sigma_\alpha\}_{\alpha \in \Omega}$ be a family of nonempty sets, $\mathbf{\Sigma} = \prod_\alpha \Sigma_\alpha$, $\beta \in \Omega$, $p \in \mathbf{\Sigma}$, $R \subseteq \mathbf{\Sigma}$, $A \in \prod_\alpha 2^{\Sigma_\alpha}$, and $B \subseteq \Sigma_\beta$. We shall make use of the following notations:

1. We denote by $\pi_\beta \colon \mathbf{\Sigma} \to \Sigma_\beta$ the $\beta-$th coordinate map, *i.e.*, $\pi_\beta(p) = p_\beta$.
2. We denote by $p[-, \beta] \colon \Sigma_\beta \to \mathbf{\Sigma}$ the map that sends $q \in \Sigma_\beta$ to the element of $\mathbf{\Sigma}$ obtained by replacing p's $\beta-$th entry by q.
3. We define $R_\beta[p] = \pi_\beta(p[\Sigma_\beta, \beta] \cap R)$. Note that $R_\beta[p] = \{q \in \Sigma_\beta \mid p[q, \beta] \in R\}$.
4. We define $A[B, \beta] \in \prod_\alpha \mathcal{L}_\alpha$ as $A[B, \beta]_\beta = B$ and $A[B, \beta]_\alpha = A_\alpha$ for $\alpha \neq \beta$.
5. We write $\overline{A} := \prod_\alpha A_\alpha$ and $\overline{A}[B, \beta] := \overline{A[B, \beta]}$.

We omit the β in $p[-,\beta]$ when no confusion can occur. For instance, we write $p[\Sigma_\beta]$ instead of $p[\Sigma_\beta,\beta]$.

Remark 2.4. $p[R_\beta[p]] = p[\Sigma_\beta] \cap R$.

Definition 2.5. Let $\{\mathcal{L}_\alpha\}_{\alpha\in\Omega}$ be a family of simple closure spaces on Σ_α. We denote by $\mathsf{S}(\mathcal{L}_\alpha, \alpha\in\Omega)$ the set all simple closure spaces $\mathcal{L}$ on $\boldsymbol{\Sigma}$ such that

- (P1) $\boldsymbol{\Sigma} = \prod_\alpha \Sigma_\alpha$,
- (P2) $\bigcup_\alpha \pi_\alpha^{-1}(a_\alpha) \in \mathcal{L}$, for all $a \in \prod_\alpha \mathcal{L}_\alpha$,
- (P3) for all $p \in \boldsymbol{\Sigma}$, $\beta \in \Omega$, and $B \subseteq \Sigma_\beta$, we have $p[B,\beta] \in \mathcal{L}$ implies $B \in \mathcal{L}_\beta$.

Let $T = \prod_\alpha T_\alpha$ with $T_\alpha \subseteq \mathsf{Aut}(\mathcal{L}_\alpha)$. We denote by $\mathcal{S}_T(\mathcal{L}_\alpha, \alpha\in\Omega)$ the set of all $\mathcal{L} \in \mathsf{S}(\mathcal{L}_\alpha, \alpha\in\Omega)$ such that

- (P4) for all $v \in T$, there is $u \in \mathsf{Aut}(\mathcal{L})$ such that $u(p)_\alpha = v_\alpha(p_\alpha)$ for all $p \in \boldsymbol{\Sigma}$ and all $\alpha \in \Omega$.

We call elements of $\mathsf{S}(\mathcal{L}_\alpha, \alpha\in\Omega)$ *weak tensor products.*

Lemma 2.6. *Let $\{\mathcal{L}_\alpha\}_{\alpha\in\Omega}$ be a family of simple closure spaces on Σ_α, $\beta\in\Omega$, and $\mathcal{L} \in \mathsf{S}(\mathcal{L}_\alpha, \alpha\in\Omega)$.*

1. *For any $a \in \prod_\alpha \mathcal{L}_\alpha$, $\overline{a} \in \mathcal{L}$.*
2. *For any $b \in \mathcal{L}_\beta$ and $p \in \prod_\alpha \Sigma_\alpha$, we have $p[b,\beta] \in \mathcal{L}$.*
3. *For any $B \subseteq \mathcal{L}_\beta$ and $a \in \prod_\alpha \mathcal{L}_\alpha$, we have $\overline{a}[\bigvee B, \beta] = \bigvee_{b\in B} \overline{a}[b,\beta]$.*

Proof. see [5] □

Definition 2.7. Let $\{\Sigma_\alpha\}_{\alpha\in\Omega}$ be a family of nonempty sets and $\{\mathcal{L}_\alpha \subseteq 2^{\Sigma_\alpha}\}_{\alpha\in\Omega}$. Let $\boldsymbol{\Sigma} = \prod_\alpha \Sigma_\alpha$. We define

$$\underset{\alpha\in\Omega}{\owedge} \mathcal{L}_\alpha := \left\{ \bigcap \omega \mid \omega \subseteq \left\{ \bigcup_\alpha \pi_\alpha^{-1}(a_\alpha) \mid a \in \prod_\alpha \mathcal{L}_\alpha \right\}, \omega \neq \emptyset \right\} ,$$

$$\underset{\alpha\in\Omega}{\ovee} \mathcal{L}_\alpha := \left\{ R \subseteq \boldsymbol{\Sigma} \mid R_\beta[p] \in \mathcal{L}_\beta, \text{ for all } p \in \boldsymbol{\Sigma},\ \beta\in\Omega \right\} ,$$

ordered by set-inclusion.

Remark 2.8. Note that $2 \owedge \mathcal{L} \cong \mathcal{L} \cong 2 \ovee \mathcal{L}$ and $|\mathsf{S}(2,\mathcal{L})| = 1$.

We end this section by recalling a definition and two results presented in [5] that we will use later.

Lemma 2.9. *Let $\{\mathcal{L}_\alpha\}_{\alpha\in\Omega}$ be a family of simple closure spaces on Σ_α and $\mathcal{L}$ a simple closure space on $\boldsymbol{\Sigma} = \prod_\alpha \Sigma_\alpha$. Suppose that Axiom P2 holds in $\mathcal{L}$. Let p, $q \in \boldsymbol{\Sigma}$.*

1. *If $p_\beta \neq q_\beta$ for at least two $\beta\in\Omega$, then $p \vee q = p \cup q$.*
2. *For all $\beta \neq \gamma \in \Omega$ and for all $b \in \mathcal{L}_\beta$ and $c \in \mathcal{L}_\gamma$ such that $p_\beta \in b$ and $p_\gamma \in c$, $p[b,\beta] \vee p[c,\gamma] = p[b,\beta] \cup p[c,\gamma]$.*

Definition 2.10. Let $\mathcal{L}$ be a simple closure space on Σ. We say that $\mathcal{L}$ is *weakly connected* if $\mathcal{L} \neq 2$ and if there is a *connected covering* of Σ, that is a family of subsets $\{A^\gamma \subseteq \Sigma \mid \gamma\in\sigma\}$ such that

1. $\Sigma = \bigcup\{A^\gamma \mid \gamma\in\sigma\}$ and $|A^\gamma| \geq 2$ for all $\gamma\in\sigma$,
2. for all $\gamma\in\sigma$ and all $p \neq q \in A^\gamma$, $p \vee q$ contains a third atom,
3. for all p, $q \in \Sigma$, there is a finite subset $\{\gamma_1, \cdots, \gamma_n\} \subseteq \sigma$ such that $p \in A^{\gamma_1}$, $q \in A^{\gamma_n}$, and such that $|A^{\gamma_i} \cap A^{\gamma_{i+1}}| \geq 2$ for all $1 \leq i \leq n-1$.

Remark 2.11. Note that weakly connected implies irreducible (see [5]). Finally, let $\mathcal{L}$ be a simple closure space. Then, if $\mathcal{L} \neq 2$ and $\mathcal{L}$ is irreducible orthocomplemented with the covering property or an irreducible DAC-lattice, then $\mathcal{L}$ is weakly connected.

Theorem 2.12. *Let Ω be a finite set and $\{\mathcal{L}_i\}_{i\in\Omega}$ a finite family of weakly connected simple closure spaces on Σ_i. Let $\mathcal{L} \in \mathsf{S}(\mathcal{L}_i, i \in \Omega)$ and $u \in \mathsf{Aut}(\mathcal{L})$. Then there is a bijection f of Ω, and for each $i \in \Omega$, there is an isomorphism $v_i \colon \mathcal{L}_i \to \mathcal{L}_{f(i)}$ such that $u(p)_{f(i)} = v_i(p_i)$ for all $p \in \mathbf{\Sigma}$ and $i \in \Omega$.*

Remark 2.13. It can be useful to have in mind the following pictures in order to read the proofs below easily. If $\Omega = \{1, 2, 3\}$, then

1. $\pi_1^{-1}(a_1) = a_1 \times \Sigma_2 \times \Sigma_3$ which we can denote by the symbol $a_1||$,
2. $\bigcup_{i=1}^{3} \pi_i^{-1}(a_i) = a_1 \times \Sigma_2 \times \Sigma_3 \cup \Sigma_1 \times a_2 \times \Sigma_3 \cup \Sigma_1 \times \Sigma_2 \times a_3$ which we can denote as $a_1|| \cup |a_2| \cup ||a_3$. Hence, we can write

$$\left(\bigcup_{i=1}^{3} \pi_i^{-1}(a_i)\right) \cap \left(\bigcup_{i=1}^{3} \pi_i^{-1}(b_i)\right) = \Big[a_1|| \cup |a_2| \cup ||a_3\Big] \cap \Big[b_1|| \cup |b_2| \cup ||b_3\Big]$$

$$= a_1 \cap b_1|| \cup a_1 b_2| \cup a_1|b_3 \cup b_1 a_2| \cup |a_2 \cap b_2| \cup |a_2 b_3 \cup b_1|a_3 \cup |b_2 a_3 \cup ||a_3 \cap b_3$$

$$= \bigcup_{f\in 3^{\{a,b\}}} \Big(\bigcap f^{-1}(1)_1\Big) \times \Big(\bigcap f^{-1}(2)_2\Big) \times \Big(\bigcap f^{-1}(3)_3\Big)$$

where a, $b \in \mathcal{L}_1 \times \mathcal{L}_2 \times \mathcal{L}_3$.

3. Sufficient and necessary conditions for $\mathcal{L}_1 \owedge \mathcal{L}_2 = \mathcal{L}_1 \ovee \mathcal{L}_2$

Let $\mathcal{L}_1, \cdots, \mathcal{L}_n$ be simple closure spaces and DAC-lattices. In this section, we prove that $\owedge_{i=1}^{n} \mathcal{L}_i = \ovee_{i=1}^{n} \mathcal{L}_i$ (*i.e.*, $\mathsf{S}(\mathcal{L}_1, \cdots, \mathcal{L}_n)$ has only one element) if and only if there is k between 1 and n such that for all $i \neq k$, $\mathcal{L}_i$ is a power set.

Definition 3.1. Let $\{\mathcal{L}_\alpha\}_{\alpha\in\Omega}$ be a family of simple closure spaces on Σ_α and $\beta \in \Omega$. Let $\mathbf{\Sigma} = \prod_\alpha \Sigma_\alpha$ and $R \subseteq \mathbf{\Sigma}$. Then, we define $\bigvee_\beta R := \bigcup_{p\in\mathbf{\Sigma}} p[\bigvee R_\beta[p]]$, where the join is being taken in $\mathcal{L}_\beta$.

Lemma 3.2. *Let $\{\mathcal{L}_\alpha\}_{\alpha\in\Omega}$ be a family of simple closure spaces on Σ_α and $\mathcal{L} \in \mathsf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$. Let $\mathbf{\Sigma} = \prod_\alpha \Sigma_\alpha$ and $R \subseteq \mathbf{\Sigma}$. Denote by $\bigvee_{\mathcal{L}}$ the join in $\mathcal{L}$.*

1. *For any $\beta \in \Omega$, $R \subseteq \bigvee_\beta R$ and $\bigvee_\beta(\bigvee_\beta R) = \bigvee_\beta R$. Moreover, if $R \subseteq S \subseteq \mathbf{\Sigma}$, then $\bigvee_\beta R \subseteq \bigvee_\beta S$.*
2. *For any $f \colon \mathbb{N} \to \Omega$ and any $n \in \mathbb{N}$, $R^n := \bigvee_{f(n)}(\cdots(\bigvee_{f(1)} R)\cdots) \subseteq \bigvee_{\mathcal{L}} R$.*
3. *$R \in \ovee_\alpha \mathcal{L}_\alpha$ if and only if $\bigvee_\beta R = R$ for all $\beta \in \Omega$.*

Proof. (1) Let $\beta \in \Omega$. First note that

$$R = R \cap \left(\bigcup_{p\in\mathbf{\Sigma}} p[\Sigma_\beta]\right) = \bigcup_{p\in\mathbf{\Sigma}} (p[\Sigma_\beta] \cap R) = \bigcup_{p\in\mathbf{\Sigma}} p[R_\beta[p]]\,.$$

As a consequence, since $R_\beta[p] \subseteq \bigvee R_\beta[p]$ in $\mathcal{L}_\beta$, we have

$$R = \bigcup_{p\in\mathbf{\Sigma}} p[R_\beta[p]] \subseteq \bigcup_{p\in\mathbf{\Sigma}} p[\bigvee R_\beta[p]] = \bigvee_\beta R,$$

hence, $R \subseteq \bigvee_\beta R$.

Further,

$$\bigvee_\beta(\bigvee_\beta R) = \bigvee_\beta \left(\bigcup_{p\in\mathbf{\Sigma}} p[\bigvee R_\beta[p]]\right) = \bigcup_{q\in\mathbf{\Sigma}} q\left[\bigvee\left(\bigcup_{p\in\mathbf{\Sigma}} p[\bigvee R_\beta[p]]\right)_\beta [q]\right]$$

$$= \bigcup_{q\in\mathbf{\Sigma}} q\left[\bigvee\left(q[\bigvee R_\beta[q]]\right)_\beta [q]\right] = \bigcup_{q\in\mathbf{\Sigma}} q\left[\bigvee\left(\bigvee R_\beta[q]\right)\right] = \bigcup_{q\in\mathbf{\Sigma}} q\left[\bigvee R_\beta[q]\right]$$

$$= \bigvee_\beta R\,.$$

Finally, let $S \subseteq \mathbf{\Sigma}$ with $R \subseteq S$, then obviously $R_\beta[p] \subseteq S_\beta[p]$ for all $p \in \mathbf{\Sigma}$, hence $\bigvee R_\beta[p] \subseteq \bigvee S_\beta[p]$. As a consequence,

$$\bigvee_\beta R = \bigcup_{p\in\mathbf{\Sigma}} p[\bigvee R_\beta[p]] \subseteq \bigcup_{p\in\mathbf{\Sigma}} p[\bigvee S_\beta[p]] = \bigvee_\beta S.$$

(2) By Axiom P3, $(\bigvee_{\mathcal{L}} R)_\beta[p] \in \mathcal{L}_\beta$ for all $p \in \mathbf{\Sigma}$ and all $\beta \in \Omega$; whence $\bigvee_\beta(\bigvee_{\mathcal{L}} R) = \bigvee_{\mathcal{L}} R$. Thus, $R^n \subseteq (\bigvee_{\mathcal{L}} R)^n = \bigvee_{\mathcal{L}} R$, since $R \subseteq \bigvee_{\mathcal{L}} R$.

(3) Suppose that $R \in \ovee_\alpha \mathcal{L}_\alpha$. Then, by Definition 2.7, $R_\beta[p] \in \mathcal{L}_\beta$ for all $p \in \mathbf{\Sigma}$ and all $\beta \in \Omega$. Hence $R_\beta[p] = \bigvee R_\beta[p]$ in $\mathcal{L}_\beta$. As a consequence, $\bigvee_\beta R = R$.

Suppose now that $\bigvee_\beta R = R$ for some $\beta \in \Omega$. Let $q \in \mathbf{\Sigma}$. Then

$$R_\beta[q] = \left(\bigvee_\beta R\right)_\beta [q] = \left(\bigcup_{p\in\mathbf{\Sigma}} p[\bigvee R_\beta[p]]\right)_\beta [q] = \left(q[\bigvee R_\beta[q]]\right)_\beta [q]$$

$$= \bigvee R_\beta[q]\,.$$

As a consequence, if $\bigvee_\beta R = R$ for all $\beta \in \Omega$, then $R_\beta[q] \in \mathcal{L}_\beta$ for all $\beta \in \mathcal{L}_\beta$ and all $q \in \mathbf{\Sigma}$, hence $R \in \ovee_\alpha \mathcal{L}_\alpha$. ☐

Remark 3.3. Note that $\bigvee_\alpha R$ is not necessarily in $\mathcal{L}$. Take for instance $\mathcal{L} = \mathcal{L}_1 \owedge \mathcal{L}_2$ and $R = \{p, q, r\}$ with $p_1 \neq r_1$, $p_2 \neq r_2$, $q_1 = p_1$, and $q_2 = r_2$. Then $\bigvee_{\owedge} R = p_1 \times (p_2 \vee r_2) \cup (p_1 \vee r_1) \times r_2$. Now, $\bigvee_2 R = p_1 \times (p_2 \vee r_2) \cup r \neq \bigvee_{\owedge} R$. But, $\bigvee_1(\bigvee_2 R) = p_1 \times (p_2 \vee r_2) \cup (p_1 \vee r_1) \times r_2 = \bigvee_{\owedge} R$.

Theorem 3.4. *Let $\{\mathcal{L}_\alpha\}_{\alpha\in\Omega}$ be a family of simple closure spaces on Σ_α and $\mathbf{\Sigma} = \prod_\alpha \Sigma_\alpha$. If there is at most one $\beta \in \Omega$ such that $\mathcal{L}_\beta \neq 2^{\Sigma_\beta}$, then $\owedge_\alpha \mathcal{L}_\alpha = \ovee_\alpha \mathcal{L}_\alpha = \{R \subseteq \mathbf{\Sigma} \mid R_\beta[p] \in \mathcal{L}_\beta, \forall p \in \mathbf{\Sigma}\}$.*

Proof. Let $R \subseteq \mathbf{\Sigma}$. Since $\owedge_\alpha \mathcal{L}_\alpha \subseteq \ovee_\alpha \mathcal{L}_\alpha$, $\bigvee_{\ovee} R \subseteq \bigvee_{\owedge} R$. We prove that $\bigvee_{\owedge} R \subseteq \bigvee_{\ovee} R$. Write $\Sigma_1 = \Sigma_\beta$, $\mathcal{L}_1 = \mathcal{L}_\beta$, and $\Sigma_2 = \prod_{\alpha\neq\beta} \Sigma_\alpha$. Hence $\mathbf{\Sigma} = \Sigma_1 \times \Sigma_2$. We denote by $\pi_i \colon \mathbf{\Sigma} \to \Sigma_i$ $(i = 1, 2)$ the $i-$th coordinate map. From definition 2.7, we find that $\owedge_{\alpha\neq\beta} \mathcal{L}_\alpha = 2^{\Sigma_2}$. Therefore, since $\owedge$ is associative, $\owedge_{\alpha\in\Omega} \mathcal{L}_\alpha = \mathcal{L}_1 \owedge 2^{\Sigma_2}$.

Let $a \in \mathcal{L}_1$ and $b \in 2^{\Sigma_2}$. Note that $R \subseteq a \times \Sigma_2 \bigcup \Sigma_1 \times b$ if and only if $g(b) := \pi_1(\Sigma_1 \times b^c \cap R) \subseteq a$, where b^c denotes the set-complement of b, *i.e.*, $b^c = \Sigma_2 \backslash b$. Note also that

$$R \subseteq (\vee g(b)) \times \Sigma_2 \cup \Sigma_1 \times (b \cap \pi_2(R))\,, \text{ for all } b \in 2^{\Sigma_2}.$$

Moreover, let $a \in \mathcal{L}_2$ and $b \in 2^{\Sigma_2}$. Then, $R \subseteq a \times \Sigma_2 \cup \Sigma_1 \times b$ implies

$$(\vee g(b)) \times \Sigma_2 \cup \Sigma_1 \times (b \cap \pi_2(R)) \subseteq a \times \Sigma_2 \cup \Sigma_1 \times b\ .$$

Finally, $g(b) = g(b \cap \pi_2(R))$ for all $b \in 2^{\Sigma_2}$. As a consequence, we find that

$$\begin{aligned}\bigvee_{\owedge} R &= \bigcap \big\{a \times \Sigma_2 \cup \Sigma_1 \times b \mid a \in \mathcal{L}_1,\, b \in 2^{\Sigma_2} \text{ and } R \subseteq a \times \Sigma_2 \cup \Sigma_1 \times b\big\} \\ &= \bigcap_{b \subseteq \pi_2(R)} \Big(\bigvee g(b)\Big) \times \Sigma_2 \cup \Sigma_1 \times b \\ &= \bigcup_{f \in 2^{\left(2^{\pi_2(R)}\right)}} \Big(\bigcap \Big\{\bigvee g(b) \mid b \in f^{-1}(1)\Big\}\Big) \times \Big(\bigcap f^{-1}(2)\Big)\ .\end{aligned}$$

Write X_f for the last term in the preceding equation and $b = \cap f^{-1}(2)$. Let $g \in 2^{\left(2^{\pi_2(R)}\right)}$ such that $b \subseteq c \Rightarrow g(c) = 2$. Then $\cap g^{-1}(2) = b$ and $X_f \subseteq X_g$. As a consequence, if for $b \subseteq \pi_2(R)$ we define $m(b) := \{c \subseteq \pi_2(R) \mid b \nsubseteq c\}$, then,

$$\bigvee_{\owedge} R = \bigcup_{b \subseteq \pi_2(R)} \Big(\bigcap \Big\{\bigvee g(c) \mid c \in m(b)\Big\}\Big) \times b.$$

Note that for all $q \in b$, $q^c \bigcap \pi_2(R) \in m(b)$. Moreover,

$$g\left(q^c \bigcap \pi_2(R)\right) = \pi_1\left(\Sigma_1 \times \left(q \bigcup \pi_2(R)^c\right) \bigcap R\right) = R_1[(\cdot, q)].$$

As a consequence,

$$\begin{aligned}\bigvee_{\owedge} R &\subseteq \bigcup_{b \subseteq \pi_2(R)} \Big(\bigcap \Big\{\bigvee R_1[(\cdot, q)] \mid q \in b\Big\}\Big) \times b \\ &= \bigcup_{q \in \pi_2(R)} \Big(\bigvee R_1[(\cdot, q)]\Big) \times q = \bigvee_{\beta} R\,.\end{aligned}$$

Finally, by Lemma 3.2, $\bigvee_\beta R = \bigvee_{\ovee} R$. As a consequence, $\bigvee_{\owedge} R \subseteq \bigvee_{\ovee} R$. □

Definition 3.5. Let Σ be a nonempty set. We denote by MO_Σ the simple closure space on Σ which contains only $\emptyset$, Σ, and all singletons of Σ. We write MO_n if $|\Sigma| = n$.

Let $\mathcal{L}$ be a simple closure space. We say that $\mathcal{L}$ *contains* MO_n if there are n atoms $p_1, \cdots, p_n$ such that $p_1 \bigvee p_n \gtrdot p_i$ for all i between 1 and n.

Let $\{\Sigma_\alpha\}_{\alpha \in \Omega}$ be a family of sets, and $\mathbf{\Sigma} = \prod_\alpha \Sigma_\alpha$. We denote by $\Xi(\mathbf{\Sigma})$ the set $\{R \subseteq \mathbf{\Sigma} \mid p_\alpha \neq q_\alpha,\, \forall p, q \in R,\, \alpha \in \Omega\}$.

Theorem 3.6. *Let $\mathcal{L}_1$ and $\mathcal{L}_2$ be simple closure spaces on Σ_1 and Σ_2 respectively. If both $\mathcal{L}_1$ and $\mathcal{L}_2$ contain MO_3, then $\mathcal{L}_1 \owedge \mathcal{L}_2 \neq \mathcal{L}_1 \ovee \mathcal{L}_2$.*

Proof. Let $R = \{p, q, r\} \in \Xi(\Sigma_1 \times \Sigma_2)$ such that for $i = 1$ and for $i = 2$, $p_i \vee q_i$ covers p_i, q_i and r_i. By Definition 2.7, $R = \{p, q, r\} \in \mathcal{L}_1 \ovee \mathcal{L}_2$. On the other hand, $R \subseteq a \times \Sigma_2 \cup \Sigma_1 \times b$ if and only if $p_1 \vee q_1 \subseteq a$ or $p_2 \vee q_2 \subseteq b$. As a consequence, $\bigvee_{\owedge} R = (p_1 \vee q_1) \times (p_2 \vee q_2) \neq R = \bigvee_{\ovee} R$. □

The next corollary is a partial converse to Theorem 3.4.

Corollary 3.7. *Let $\{\mathcal{L}_i\}_{1\leq i\leq n}$ be a finite family of simple closure spaces on Σ_i. Suppose that each $\mathcal{L}_i$ is a DAC-lattice. If $\owedge_{i=1}^n\mathcal{L}_i = \ovee_{i=1}^n\mathcal{L}_i$, then there is at most one i such that $\mathcal{L}_i \neq 2^{\Sigma_i}$.*

Proof. Suppose that $\mathcal{L}_i \neq 2^{\Sigma_i}$ for $i = m$ and for $i = k$ with $m \neq k$ between 1 and n, and that $\owedge_{i=1}^n\mathcal{L}_i = \ovee_{i=1}^n\mathcal{L}_i$. Then, for $i = m$ and for $i = k$, there is an atom p_i which is not a central element. Therefore, there is an atom q_i such that $p_i \bigvee q_i$ contains a third atom (see [6], Theorems 28.8, 27.6 and Lemma 11.6).

Let $\mathcal{L}_0 := \owedge_{i\neq m}\mathcal{L}_i$ and $\mathcal{L}_1 := \ovee_{i\neq m}\mathcal{L}_i$. By hypothesis, $\mathcal{L}_m \owedge \mathcal{L}_0 = \mathcal{L}_m \ovee \mathcal{L}_1$. Therefore, from Lemma 2.6 and Axiom P3, $\mathcal{L}_0 = \mathcal{L}_1$. Let $r \in \prod_{i\neq m}\Sigma_i$ be an atom of $\mathcal{L}_0$. From Lemma 2.6, $r[p_k] \vee_{\owedge} r[q_k] = r[p_k \vee q_k]$, hence contains a third atom, therefore $\mathcal{L}_0$ contains MO_3. As a consequence, from Theorem 3.6, we find that

$$\mathcal{L}_m \ovee \mathcal{L}_1 = \mathcal{L}_m \owedge \mathcal{L}_0 \neq \mathcal{L}_m \ovee \mathcal{L}_0 = \mathcal{L}_m \ovee \mathcal{L}_1,$$

a contradiction. □

Theorem 3.8. *Let $\mathcal{L}$ be a coatomistic simple closure space on Σ. Suppose that for any countable set A of coatoms $\mathcal{L}$, $\bigcup A \neq \Sigma$. For all integers i, let $\mathcal{L}_i = \mathcal{L}$. Then $\owedge_{i=1}^\infty\mathcal{L}_i \neq \ovee_{i=1}^\infty\mathcal{L}_i$.*

Proof. For all i, let $\Sigma_i = \Sigma$. Denote $\prod_{i=1}^\infty \Sigma_i$ by $\mathbf{\Sigma}$. Let

$$R = \{p \in \mathbf{\Sigma} \mid p_n = p_m,\ \forall m, n\} \neq \mathbf{\Sigma}.$$

From Definition 2.7, $R \in \ovee_{i=1}^\infty\mathcal{L}_i$. On the other hand, by hypothesis, for any $a \in \prod_{i=1}^\infty(\mathcal{L}_i\backslash\{1\})$, $R \nsubseteq \bigcup_{i=1}^\infty \pi_i^{-1}(a_i)$, hence $\bigvee_{\owedge} R = 1$. □

Example 3.9. The Hypothesis of Theorem 3.8 is fulfilled for instance if $\mathcal{L} = \mathsf{P}(\mathcal{H})$ with $\mathcal{H}$ a real or complex Hilbert space.

4. Orthocomplementation

This section is devoted to our main results, which show, subject to weak conditions, that if $\mathcal{L} \in \mathsf{S}(\mathcal{L}_1, \cdots, \mathcal{L}_n)$ is orthocomplemented, then $\mathcal{L} = \owedge_{i=1}^n\mathcal{L}_i$. Using Theorem 2.12, we prove that this holds if each $\mathcal{L}_i$ is weakly connected (hence, irreducible) and coatomistic and if $\mathcal{L}$ is moreover transitive. For the second result, we assume that each $\mathcal{L}_i$ is orthocomplemented and that all its irreducible components different from 2 are weakly connected, but we do not need to assume that $\mathcal{L}$ is transitive.

Definition 4.1. Let $\{\Sigma_\alpha\}_{\alpha\in\Omega}$ be a family of nonempty sets, $p \in \mathbf{\Sigma} = \prod_\alpha \Sigma_\alpha$, β, $\gamma \in \Omega$, $A_\beta \subseteq \Sigma_\beta$, and $A_\gamma \subseteq \Sigma_\gamma$. Then

$$p[A_\beta, A_\gamma] := \{q \in \mathbf{\Sigma} \mid q_\beta \in A_\beta,\ q_\gamma \in A_\gamma,\ q_\alpha = p_\alpha,\ \forall \alpha \neq \beta,\ \gamma\}.$$

Lemma 4.2. *Let $\{\mathcal{L}_i\}_{1\leq i\leq n}$ be a finite family of simple closure spaces on Σ_i, $x \in \prod_{i=1}^n \mathcal{L}_i$ with x_i coatoms, and let $\mathcal{L} \in \mathsf{S}(\mathcal{L}_i, 1 \leq i \leq n)$. Then:*

1. *$X := \bigcup_{i=1}^n \pi_i^{-1}(x_i)$ is a coatom of $\mathcal{L}$.*
2. *Let $a_j \in \mathcal{L}_j$ for some j between 1 and n and let Z be a coatom of $\mathcal{L}$ above $\bigcup_{i\neq j} \pi_i^{-1}(x_i) \bigcup \pi_j^{-1}(a_j)$. Then there is a coatom z_j of $\mathcal{L}_j$ such that $Z = \bigcup_{i\neq j} \pi_i^{-1}(x_i) \bigcup \pi_j^{-1}(z_j)$.*

Proof. (1): By Axiom P2, $X \in \mathcal{L}$. Let $\mathbf{\Sigma} = \prod_{i=1}^{n} \Sigma_i$ and $p \in \mathbf{\Sigma}$ not in X. Write $R^0 := p \cup X$ and $y := p \vee X$. Define $R^N := \bigvee_n \cdots \bigvee_1 R^{N-1}$. By Lemma 3.2, $R^N \subseteq y$, for all N. Now,

$$R^0_k[q] = \{s \in \Sigma_k \mid q[s,k] \in p \cup X\} = \{p\}_k[q] \cup X_k[q];$$

therefore,

$$q[R^0_k[q]] = q[X_k[q]] \text{ or } q[R^0_k[q]] = p[x_k \cup p_k],$$

hence

$$q[\bigvee R^0_k[q]] = q[\vee X_k[q]] \subseteq X \text{ or } q[\vee R^0_k[q]] = p[\Sigma_k]\,.$$

As a consequence, $\bigvee_k R^0 = p[\Sigma_k] \cup R^0$, thus $R^1 = R^0 \bigcup_{k=1}^{n} p[\Sigma_k]$. Further,

$$R^2 = R^0 \bigcup \{p[\Sigma_{k_1}, \Sigma_{k_2}] \mid 1 \leq k_1 \leq n-1 \text{ and } k_1 + 1 \leq k_2 \leq n\}\,.$$

Hence obviously, for $N = n$, $R^N = \mathbf{\Sigma}$.

(2): Write $Y := \bigcup_{i \neq j} \pi_i^{-1}(x_i) \bigcup \pi_j^{-1}(a_j)$. Let $r \in Z$ not in Y. Note that $r[a_j] \subseteq Y$. Therefore, by Lemma 2.6, $r[a_j \bigvee r_j] \subseteq Z$. Let $k \neq j$ between 1 and n. Then, $r[x_k, a_j \vee r_j] \subseteq Y$. Hence, since $r[a_j \vee r_j] \subseteq Z$, by Lemma 2.6, $r[\Sigma_k, a_j \vee r_j] \subseteq Z$. Repeating this argument, we find that $\pi_j^{-1}(a_j \vee r_j) \subseteq Z$. As a consequence,

$$Z = Y \bigvee_{r \in Z \setminus Y} r \subseteq \bigcup_{i \neq j} \pi_i^{-1}(x_i) \bigvee_{r \in Z \setminus Y} \pi_j^{-1}(a_j \vee r_j) \subseteq Z$$

which proves that $Z = \bigcup_{i \neq j} \pi_i^{-1}(x_i) \cup \pi_j^{-1}(z_j)$ for some coatom z_j of $\mathcal{L}_j$. □

Proposition 4.3. *Let $\{\mathcal{L}_\alpha\}_{\alpha \in \Omega}$ be a family of simple closure spaces on Σ_α.*

1. *If $\owedge_\alpha \mathcal{L}_\alpha$ is coatomistic, then all $\mathcal{L}_\alpha$'s are coatomistic.*
2. *Suppose that Ω is finite and that all $\mathcal{L}_\alpha$'s are weakly connected. If $\owedge_\alpha \mathcal{L}_\alpha$ is orthocomplemented, then all $\mathcal{L}_\alpha$'s are orthocomplemented.*

Proof. (1): Let $\mathbf{\Sigma} = \prod_\alpha \Sigma_\alpha$ and $p \in \mathbf{\Sigma}$. From Definition 2.7, a coatom of $\owedge_\alpha \mathcal{L}_\alpha$ over p is necessarily of the form $\bigcup_\alpha \pi_\alpha^{-1}(x_\alpha)$ where all x_α's are coatoms and with $p_\beta \in x_\beta$ for some $\beta \in \Omega$. Let $\Sigma'_\alpha[p_\alpha]$ denote the set of coatoms of $\mathcal{L}_\alpha$ above p_α (write $\Sigma'_\alpha := \Sigma'_\alpha[0_\alpha]$). Define $\chi = \{x \in \prod_\alpha \Sigma'_\alpha \mid p_\beta \in x_\beta \text{ for some } \beta \in \Omega\}$. If $\owedge_\alpha \mathcal{L}_\alpha$ is coatomistic, then

$$p = \bigcap \left\{ \bigcup_\alpha \pi_\alpha^{-1}(x_\alpha) \mid x \in \ \chi \right\} = \bigcup_{f \in \Omega^\chi} \left(\prod_{\alpha \in \Omega} \left(\bigcap f^{-1}(\alpha)_\alpha \right) \right).$$

Hence, if $\Sigma'_{\alpha_0}[p_{\alpha_0}] = \emptyset$ for some $\alpha_0 \in \Omega$, then $\pi_{\alpha_0}(p) = \Sigma_{\alpha_0}$ which means that $\mathcal{L}_{\alpha_0} = 2$. As a consequence, all $\mathcal{L}_\alpha$'s are coatomic and we can assume that $\Sigma'_\alpha[p_\alpha] \neq \emptyset$ for all $\alpha \in \Omega$. Moreover, it follows that $p_\alpha = \bigcap \{x_\alpha \in \Sigma'_\alpha[p_\alpha]\}$, for all $\alpha \in \Omega$, hence that all $\mathcal{L}_\alpha$'s are coatomistic.

(2): Let $\beta \in \Omega$ and $a \in \mathcal{L}_\beta$. We first prove that there is $b \in \mathcal{L}_\beta$ such that $\pi_\beta^{-1}(a)' = \pi_\beta^{-1}(b)$. Let $f \colon \mathbf{\Sigma} \to \prod \Sigma'_\alpha$ such that $p' = \bigcup_\alpha \pi_\alpha^{-1}(f(p)_\alpha)$ for all $p \in \mathbf{\Sigma}$. Note that f is injective.

(2.1) **Claim**: Let $p, q \in \mathbf{\Sigma}$. If $f(p)_\alpha \neq f(q)_\alpha$ for at least two $\alpha \in \Omega$, then $p \vee q = p \cup q$.



FIGURE 1.

Proof: Let $\Omega_{\neq} = \{\alpha \in \Omega \mid f(p)_\alpha \neq f(q)_\alpha\}$ and $\gamma \neq \delta \in \Omega_{\neq}$. Let $r \in p \vee q$. Then,

$$\bigcup_\alpha \pi_\alpha^{-1}(f(r)_\alpha) = r' \supseteq p' \bigcap q' \supseteq$$
$$\left(\pi_\gamma^{-1}(f(p)_\gamma) \cap \pi_\delta^{-1}(f(q)_\delta)\right) \cup \left(\pi_\delta^{-1}(f(p)_\delta) \cap \pi_\gamma^{-1}(f(q)_\gamma)\right) .$$

Using the notation of Remark 2.13, we can write for $\Omega = \{1, 2, 3\}$, $\gamma = 1$ and $\delta = 2$,

$$f(r)_1|| \cup |f(r)_2| \cup ||f(r)_3 \supseteq f(p)_1 f(q)_2| \cup f(q)_1 f(p)_2|$$

Therefore, $f(r)_\gamma = x_\gamma$ and $f(r)_\delta = x_\delta$ with $x = f(p)$ or $x = f(q)$. As a consequence, $r' = p'$ or $r' = q'$, hence $r = p$ or $r = q$, proving the claim.

(2.2) **Claim**: Let p, $q \in \mathbf{\Sigma}$ such that p and q differ only by one component, say γ. Then $f(p)_\alpha = f(q)_\alpha$ for all $\alpha \neq \gamma$.

Proof: Since $\mathcal{L}_\gamma$ is weakly connected, from part 2.1 we find that there is $\delta \in \Omega$ such that for all $r \in \Sigma_\gamma$; we have $f(p[r,\gamma])_\alpha = f(p)_\alpha$ for all $\alpha \neq \delta$ (see Figure 1). Hence,

$$\begin{aligned} p[\Sigma_\gamma]' &= \bigcap_{r \in \Sigma_\gamma} \left(\bigcup_{\alpha \neq \delta} \pi_\alpha^{-1}(f(p)_\alpha) \cup \pi_\delta^{-1}(f(p[r,\gamma]_\delta)\right) \\ &= \bigcup_{\alpha \neq \delta} \pi_\alpha^{-1}(f(p)_\alpha) \cup \pi_\delta^{-1}(x) , \end{aligned}$$

for some $x \in \mathcal{L}_\delta$. Now $p[\Sigma_\gamma] \bigcap \pi_\gamma^{-1}(f(p)_\gamma) \neq 0$. Therefore, if $\delta \neq \gamma$, then $p[\Sigma_\gamma] \bigcap p[\Sigma_\gamma]' \neq 0$, a contradiction. As a consequence, $\delta = \gamma$ and $x = 0$, proving the claim.

Let p, $q \in \mathbf{\Sigma}$ such that $p \in \pi_\beta^{-1}(a)'$, for some $a \in \mathcal{L}_\beta$, and such that p and q differ only by one component, say γ, with $\gamma \neq \beta$. Then $\pi_\beta^{-1}(a) \subseteq p' = \bigcup_\alpha \pi_\alpha^{-1}(f(p)_\alpha)$, hence $a \subseteq f(p)_\beta$. Now, from part 2.2, $f(q)_\beta = f(p)_\beta$, therefore $\pi_\beta^{-1}(a) \subseteq q'$, thus $q \in \pi_\beta^{-1}(a)'$. As a consequence, for all $q \in \mathbf{\Sigma}$ such that $q_\beta = p_\beta$, $q \in \pi_\beta^{-1}(a)'$.

Thus, we have proved that there is an element in $\mathcal{L}_\beta$, which we denote by $a^{\perp_\beta}$, such that $\pi_\beta^{-1}(a)' = \pi_\beta^{-1}(a^{\perp_\beta})$. Obviously, the mapping $\perp_\beta \colon \mathcal{L}_\beta \to \mathcal{L}_\beta$ is an orthocomplementation. □

Theorem 4.4. *Let* $\{\mathcal{L}_i\}_{1 \leq i \leq n}$ *be a finite family of coatomistic weakly connected simple closure spaces on* Σ_i*, and let* $\mathcal{L} \in \mathsf{S}(\mathcal{L}_i, 1 \leq i \leq n)$*. If* $\mathcal{L}$ *is transitive and orthocomplemented, then* $\mathcal{L} = \textcircled{\scriptsize{\wedge}}_{i=1}^n \mathcal{L}_i$*.*

Proof. Let $\mathbf{\Sigma} = \prod_{i=1}^n \Sigma_i$. We denote the orthocomplementation of $\mathcal{L}$ by $'$. Let $x \in \prod_{i=1}^n \mathcal{L}_i$ with x_i coatoms, and $X := \bigcup_{i=1}^n \pi_i^{-1}(x_i)$. By Lemma 4.2, $X' = p$, for some $p \in \mathbf{\Sigma}$. Let $q \in \mathbf{\Sigma}$. Since $\mathcal{L}$ is transitive, there is $u \in \mathsf{Aut}(\mathcal{L})$ such that $u(p) = q$, hence $q' = u(p)'$. Define $u' \in \mathsf{Aut}(\mathcal{L})$ as $u'(a) := u(a')'$. Then $q' = u(p)' = u'(p') = u'(X)$. From Theorem 2.12, u' factors, therefore there is $y \in \prod_{i=1}^n \mathcal{L}_i$ with y_i coatoms such that $q' = Y := \bigcup_{i=1}^n \pi_i^{-1}(y_i)$. $\square$

Remark 4.5. If $\mathcal{L} \in \mathcal{S}_T(\mathcal{L}_i, 1 \leq i \leq n)$, $\mathcal{L}_i$ are transitive, and the u in Axiom P4 is an ortho-isomorphism of $\mathcal{L}$ for all $v_i \in T_i$, then the proof does not require Theorem 2.12. Below we give a second proof which requires neither Theorem 2.12 nor that $\mathcal{L}$ be transitive.

Definition 4.6. Let $\{\mathcal{L}_\alpha\}_{\alpha\in\Omega}$ be a family of orthocomplemented simple closure spaces on Σ_i. For $p \in \prod_{\alpha\in\Omega} \Sigma_\alpha$ define $p^{\#} = \bigcup_{\alpha\in\Omega} \pi_\alpha^{-1}(p_\alpha^{\perp_\alpha})$

Remark 4.7. It is easy to check that the mapping $p \mapsto p^{\#}$ is an orthocomplementation of $\owedge_{\alpha\in\Omega}\mathcal{L}_\alpha$ (see [5] for details).

Remark 4.8. Let $\mathcal{L}$ be an orthocomplemented simple closure space on Σ. Let $\mathcal{Z}(\mathcal{L})$ denotes the center of $\mathcal{L}$. Then $\mathcal{Z}(\mathcal{L}) = \{a \in \mathcal{L} \mid a^\perp = a^c\}$, where $a^c := \Sigma\backslash a$. For $p \in \Sigma$, we write $e(p)$ for the central cover of p, that is $e(p) = \bigcap\{a \in \mathcal{Z}(\mathcal{L}) \mid p \in a\}$.

Theorem 4.9. *Let $\{\mathcal{L}_i\}_{1\leq i\leq n}$ be a finite family of orthocomplemented simple closure spaces on Σ_i. Suppose moreover that for each i, all irreducible components of $\mathcal{L}_i$ different from 2 are weakly connected. Let $\mathcal{L} \in \mathsf{S}(\mathcal{L}_i, 1 \leq i \leq n)$. If $\mathcal{L}$ is orthocomplemented, then $\mathcal{L} = \owedge_{i=1}^n \mathcal{L}_i$.*

Proof. Let $\mathbf{\Sigma} = \prod_{i=1}^n \Sigma_i$. We denote the orthocomplementation of $\mathcal{L}$ by $'$ and the orthocomplementation of $\mathcal{L}_i$ by $\perp_i$. Since coatoms of $\owedge_i\mathcal{L}_i$ are coatoms of $\mathcal{L}$, we can define a map ϕ on $\mathbf{\Sigma}$ as $\phi(p) = p^{\#'}$. Note that ϕ is injective. We prove in four steps that ϕ is surjective.

(1) **Claim**: Let $p \in \mathbf{\Sigma}$ and $a_j \in \mathcal{L}_j$ for some j between 1 and n; then $\phi(p[a_j]) = p[a_j]^{\#'}$.

Proof: If $q \in p[a_j]$, then $p[a_j]^{\#} \subseteq q^{\#}$, hence $\phi(q) \in p[a_j]^{\#'}$. On the other hand, if $q \in p[a_j]^{\#'}$, then, $p[a_j]^{\#} = \bigcup_{i\neq j} \pi_i^{-1}(p_i^{\perp_i}) \cup \pi_j^{-1}(a_j^{\perp_j}) \subseteq q'$.

Thus, by Lemma 4.2, there is an atom s_j with $a_j^{\perp_j} \subseteq s_j^{\perp_j}$ and

$$q' = \bigcup_{i\neq j} \pi_i^{-1}(p_i^{\perp_i}) \bigcup \pi_j^{-1}(s_j^{\perp_j}).$$

Therefore, $q = p[s_j]^{\#'}$ with $s_j \in a_j$. Hence $q \in \phi(p[a_j])$, proving the claim.

(2) **Claim**: Let $p \in \mathbf{\Sigma}$. For all j between 1 and n there is $q \in \prod_{i=1}^n e(p_i)$ and k between 1 and n such that $\phi(p[e(p_j)]) \subseteq q[e(p_k)]$.

Proof: Note that if $a \in \mathcal{L}$, then $a' \subseteq a^c$, where a^c denotes the set-complement of a, *i.e.*, $a^c = \mathbf{\Sigma}\backslash a$. By Claim 1,

$$\phi(p[e(p_j)]) \subseteq (p[e(p_j)]^{\#})^c = e(p_j)\textstyle\prod_{i\neq j}((p_i^{\perp_i})^c) \subseteq \prod_{i=1}^n e(p_i),$$

since if $q_i \in e(p_i)$, then $(q_i^{\perp_i})^c \subseteq e(p_i)$. If $[0, e(p_j)] = 2$ (*i.e.*, $e(p_j) = p_j$), then the proof of Claim 2 is finished.

Otherwise, let $t_j \neq s_j \in e(p_j)$. Then,

$$\begin{aligned} q \in \phi(p[t_j]) \vee \phi(p[s_j]) &\text{ iff } p[t_j]^{\#} \cap p[s_j]^{\#} \subseteq q' \\ &\text{ iff } \bigcup_{i\neq j} \pi_i^{-1}(p_i^{\perp_i}) \cup \pi_j^{-1}((t_j \bigvee s_j)^{\perp_j}) \subseteq q'\,, \end{aligned}$$

hence, by Lemma 4.2, if and only if $q = \phi(p[r_j])$ for some $r_j \in t_j \vee s_j$.

As a consequence, if $t_j \bigvee s_j$ contains a third atom, so does $\phi(p[t_j]) \vee \phi(p[s_j])$. Hence, by Lemma 2.9 part 1, $\phi(p[t_j])$ and $\phi(p[s_j])$ differ only by one component, say k. Therefore, since $\mathcal{L}_j$ is weakly connected, for all A_j^γ in the connected covering of Σ_j, there is $q^\gamma \in \prod_{i=1}^n e(p_i)$ and k_γ such that $\phi(p[e(p_j) \bigcap A_j^\gamma]) \subseteq q^\gamma[e(p_{k_\gamma})]$. From Hypotheses 1 and 3 in Definition 2.10, the maps $\gamma \mapsto k_\gamma$ and $\gamma \mapsto q_\gamma$ are constant since ϕ is injective. This completes the proof of the claim.

(3) **Claim**: For all $p \in \mathbf{\Sigma}$ and all j between 1 and n, there is $q \in \prod_{i=1}^n e(p_i)$ such that $\phi(p[e(p_j)]) = q[e(p_j)]$.

Proof: From Claim 2, there is $q \in \prod_{i=1}^n e(p_i)$, k between 1 and n, and $b_k \subseteq e(p_k)$, such that $\phi(p[e(p_j)]) = q[b_k]$.

Assume first that $k \neq j$. Let $R^0 := q[b_k] \cup p[e(p_j)]^{\#}$. By part 1,

$$R^{0\prime} = q[b_k]' \cap p[e(p_j)]^{\#\prime} = \phi(p[e(p_j)])' \cap \phi(p[e(p_j)]) = 0,$$

hence $\bigvee R^0 = 1$. On the other hand, note that since $q_j \in e(p_j)$, $q_j \vee e(p_j)^c = q_j \cup e(p_j)^c$. If $e(p_j) = p_j$, then part 3 is trivial. Hence we can assume that $e(p_j) \neq p_j$, thus that $q_j \vee e(p_j)^c \neq 1$, in other words that $e(p_j) \cap q_j^{\perp_j} \neq \emptyset$. Now, for any $r_j \in e(p_j) \cap q_j^{\perp_j}$, we have

$$R^0 = q[b_k] \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp_i}) \cup \pi_j^{-1}(e(p_j)^c) \subseteq \pi_j^{-1}(r_j^{\perp_j}) \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp_i}),$$

whence by Axiom P2, $\bigvee R^0 \neq 1$, a contradiction. As a consequence, $k = j$.

Let $R := q[b_j] \bigcup p[e(p_j)]^{\#}$. From Claim 1, $R' = \phi(p[e(p_j)])' \cap \phi(p[e(p_j)]) = 0$. Therefore, $\bigvee R = 1$. Now, $R \subseteq \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp_i}) \cup \pi_j^{-1}(e(p_j)^c \vee b_j)$. Note that for all $a \in \mathcal{L}_i$ with $a \subseteq e(p_j)$, we have $e(p_j) \cap (a \vee e(p_j)^{\perp_j}) = a$. Therefore, $a \vee e(p_j)^{\perp_j} = a \cup e(p_j)^{\perp_j}$. Hence, we find that

$$R \subseteq \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp_i}) \cup \pi_j^{-1}(e(p_j)^c \cup b_j).$$

Since $\bigvee R = 1$, from Axiom P2 we find that $b_j = e(p_j)$, proving the claim.

(4): Let $p \in \mathbf{\Sigma}$ and $s \in \prod_{i=1}^n e(p_i)$. By Claim 3, $\phi(p[e(p_1)]) = q^1[e(p_1)]$. Therefore, there is $r_1 \in e(p_1)$ such that $\phi(p[r_1])_1 = s_1$. Let $k \leq n$ and $r_1 \in e(p_1), \cdots, r_k \in e(p_k)$ such that $\phi(p[r_1, \cdots, r_k])_i = s_i$, for all $i \leq k$, and such that $\phi(p[r_1, \cdots, r_k])_{k+1}$ is different from s_{k+1}. By Claim 3,

$$\phi(p[r_1, \cdots, r_k, e(p_{k+1})]) = q^{k+1}[s_1, \cdots s_k, e(p_{k+1})].$$

Hence there is $r_{k+1} \in e(p_{k+1})$ such that $\phi(p[r_1, \cdots, r_{k+1}])_i = s_i$, for all i between 1 and $k+1$. As a consequence, ϕ is surjective. $\square$

5. COVERING PROPERTY

In this Section, we prove, under some assumptions, that the top element $\ovee_\alpha \mathcal{L}_\alpha$ has the covering property if and only at most one $\mathcal{L}_\alpha$ is not a power set. We reproduce the analogue result concerning the bottom element $\owedge_\alpha \mathcal{L}_\alpha$ which is due to Aerts [1]. Moreover, for $\mathcal{L}_i = \mathsf{MO}_{\Sigma_i}$ $(i = 1, 2)$ and $T = \mathsf{Aut}(\mathcal{L}_1) \times \mathsf{Aut}(\mathcal{L}_2)$, we prove that there is a unique $\mathcal{L} \in \mathcal{S}_T(\mathcal{L}_1, \mathcal{L}_2)$ with the covering property.

Theorem 5.1 (D. Aerts, [1]). *Let $\{\mathcal{L}_\alpha\}_{\alpha \in \Omega}$ be a family of orthocomplemented simple closure spaces on Σ_α. If $\owedge_\alpha \mathcal{L}_\alpha$ has the covering property or is orthomodular, then there is at most one $\beta \in \Omega$ such that $\mathcal{L}_\beta \neq 2^{\Sigma_\beta}$.*

Proof. Let $\mathcal{L}$ be an orthocomplemented simple closure space on Σ and let p, $q \in \Sigma$ such that $p \vee q = p \cup q$. Define $x := q^{\perp} \cap (p \vee q)$, then $x = 0$ or $x = p$. If $\mathcal{L}$ has the covering property, then $1 \neq x^{\perp}$, whereas if $\mathcal{L}$ is orthomodular, $x \vee q = p \vee q$. As a consequence, $x = p$, hence $p \perp q$.

Let $\beta \in \Omega$. Suppose that $\mathcal{L}_\beta \neq 2^{\Sigma_\beta}$. Then there are two non orthogonal atoms, say r_β and s_β. Let r_γ, $s_\gamma \in \Sigma_\gamma$ for some $\gamma \in \Omega$ different from β, and let p, $q \in \prod_\alpha \Sigma_\alpha$ defined as $p_\alpha = q_\alpha$, for all $\alpha \neq \beta$, γ, and $p_\alpha = r_\alpha$ and $q_\alpha = s_\alpha$ if $\alpha = \beta$ or γ. By Lemma 2.9, $p \vee q = p \cup q$. Therefore, by what precedes, since $\#$ is an orthocomplementation of $\owedge_\alpha \mathcal{L}_\alpha$, so $p \# q$, hence by Definition 4.6, $r_\gamma \perp_\gamma s_\gamma$. As a consequence, $\mathcal{L}_\gamma = 2^{\Sigma_\gamma}$. □

Proposition 5.2. *Let $\{\mathcal{L}_\alpha\}_{\alpha \in \Omega}$ be a family of simple closure spaces on Σ_α. If $\ovee_\alpha \mathcal{L}_\alpha$ has the covering property, then all $\mathcal{L}_\alpha$'s have the covering property.*

Proof. Let $a_\beta \in \mathcal{L}_\beta$, $q_\beta \in \Sigma_\beta$ not in a_β, and $p \in \prod_\alpha \Sigma_\alpha$. By the covering property, we find that $p[q_\beta] \bigvee p[a_\beta] \gtrdot p[a_\beta]$; Whence by Lemma 2.6.3, $q_\beta \bigvee a_\beta \gtrdot a_\beta$. □

Remark 5.3. In the next theorem, we assume that each α, $\mathcal{L}_\alpha$ is a simple closure space on Σ_α, that $\mathcal{L}_\alpha \neq 2^{\Sigma_\alpha}$, and that $\mathcal{L}_\alpha$ contains MO_4. This is for instance the case if $\mathcal{L}_\alpha$ is orthocomplemented orthomodular with the covering property. Indeed, if $\mathcal{L}_\alpha \neq 2^{\Sigma_\alpha}$, there is an atom p_α which is not central, hence such that $e(p_\alpha)$ contains at least two atoms, say r and s. Moreover, $r \vee s$ contains at least three atoms, for $\mathcal{L}_\alpha$ has the covering property. Finally, since $\mathcal{L}_\alpha$ is orthomodular, $[0, r \vee s]$ is orthocomplemented, hence contains at least four atoms.

Theorem 5.4. *Let $\{\mathcal{L}_\alpha\}_{\alpha \in \Omega}$ be a family of simple closure spaces on Σ_α.*

1. *If each $\mathcal{L}_\alpha$ has the covering property and if there is at most one $\beta \in \Omega$ such that $\mathcal{L}_\beta \neq 2^{\Sigma_\beta}$, then $\ovee_\alpha \mathcal{L}_\alpha$ has the covering property.*
2. *Suppose that each $\mathcal{L}_\alpha$ different from 2^{Σ_α} contains MO_4. If $\ovee_\alpha \mathcal{L}_\alpha$ has the covering property, then there is at most one $\beta \in \Omega$ such that $\mathcal{L}_\beta \neq 2^{\Sigma_\beta}$.*

Proof. (1): Let $\mathbf{\Sigma} = \prod_\alpha \Sigma_\alpha$, $a \in \ovee_\alpha \mathcal{L}_\alpha$, $q \in \mathbf{\Sigma}$ not in a, and $R = q \cup a$. By Lemma 3.2, $q \vee_{\ovee} a = \bigvee_\beta R = \bigcup_{p \in \mathbf{\Sigma}} p[\bigvee R_\beta[p]]$. Now, by definition,

$$R_\beta[p] = \pi_\beta(p[\Sigma_\beta] \cap (q \cup a)) = q_\beta[p] \cup a_\beta[p],$$

and $q_\beta[p] = q_\beta$ if $p \in q[\Sigma_\beta]$, and $q_\beta[p] = \emptyset$ otherwise. Hence, for all p not in $q[\Sigma_\beta]$, $R_\beta[p] = a_\beta[p] \in \mathcal{L}_\beta$, and if $p \in q[\Sigma_\beta]$, then $R_\beta[p] = q_\beta \cup a_\beta[p]$. Therefore, we find that

$$q \cup a \subseteq q \vee_{\ovee} a = q[q_\beta \vee a_\beta[q]] \bigcup_{p \notin q[\Sigma_\beta]} p[a_\beta[p]] \subseteq a \cup q[q_\beta \bigvee a_\beta[q]].$$

As a consequence, $q \bigvee_{\ovee} a \gtrdot a$.

(2): Let $\beta \neq \gamma \in \Omega$. Suppose that neither $\mathcal{L}_\beta = 2^{\Sigma_\beta}$ nor $\mathcal{L}_\gamma = 2^{\Sigma_\gamma}$. Let p, q, r, s, $t \in \mathbf{\Sigma}$ such that $p_\alpha = q_\alpha = r_\alpha = s_\alpha = t_\alpha$, for all α different from β and γ, and such that $t_\beta = p_\beta$ and $t_\gamma = q_\gamma$. Assume moreover that for $\alpha = \beta$ and for $\alpha = \gamma$, p_α, q_α, r_α and s_α are all different and that $p_\alpha \bigvee q_\alpha$ covers p_α, q_α, r_α and s_α. By Definition 2.7, $a = \{p, q, r\}$ and $b = \{p, q, r, s\}$ are in $\ovee_\alpha \mathcal{L}_\alpha$. Let $R^0 = a \bigcup t$.

Then (see Lemma 3.2),

$$\begin{aligned} R^1 &:= \bigvee_{\gamma} R^0 = R^0 \cup p[p_\gamma \vee q_\gamma] \\ R^2 &:= \bigvee_{\beta} R^1 = R^0 \cup p[p_\gamma \vee q_\gamma] \cup q[p_\beta \vee q_\beta] \cup r[p_\beta \vee q_\beta] \\ R^3 &:= \bigvee_{\gamma} R^2 = p[p_\beta \vee q_\beta, p_\gamma \vee q_\gamma]\,. \end{aligned}$$

Hence, by Lemma 3.2, $a \bigvee_{\ovee} t = R^3 \supsetneq b \supsetneq a$, therefore $\ovee_\alpha \mathcal{L}_\alpha$ does not have the covering property. □

Definition 5.5. Let Σ_1, Σ_2 be sets, $\mathcal{L}_1 = \mathsf{MO}_{\Sigma_1}$, and $\mathcal{L}_2 = \mathsf{MO}_{\Sigma_2}$ (see Definition 3.5). Then,

$$\mathcal{L}_1 \circ \mathcal{L}_2 := \mathcal{L}_1 \owedge \mathcal{L}_2 \cup \{R \in \Xi(\Sigma_1 \times \Sigma_2) \mid |R| = 3\}.$$

Theorem 5.6. *Let Σ_1 and Σ_2 be sets, $\mathcal{L}_1 = \mathsf{MO}_{\Sigma_1}$, $\mathcal{L}_2 = \mathsf{MO}_{\Sigma_2}$, $T = \mathsf{Aut}(\mathcal{L}_1) \times \mathsf{Aut}(\mathcal{L}_2)$, and $\mathcal{L} \in \mathcal{S}_T(\mathcal{L}_1, \mathcal{L}_2)$. Suppose that for $i = 1$ and 2, $|\Sigma_i| = 3$ or $|\Sigma_i| \geq 5$. Then $\mathcal{L}$ has the covering property if and only if $\mathcal{L} = \mathcal{L}_1 \circ \mathcal{L}_2$.*

Proof. ($\Leftarrow$): Let $\mathbf{\Sigma} = \Sigma_1 \times \Sigma_2$, $\Xi = \Xi(\Sigma_1 \times \Sigma_2)$, and $a \in \mathcal{L}_1 \circ \mathcal{L}_2$. Then $a \in \Xi$ and $|a| = 3$, or $a \in \mathcal{L}_1 \owedge \mathcal{L}_2$. Hence one of the following cases holds.

1. $a \in \mathbf{\Sigma}$.
2. $a \in \Xi$ and $|a| = 2$ or 3.
3. $a = p_1 \times \Sigma_2$ or $a = \Sigma_2 \times p_2$ for some $p = (p_1, p_2) \in \mathbf{\Sigma}$.
4. $a = p_1 \times \Sigma_2 \cup \Sigma_1 \times p_2$, for some $p \in \mathbf{\Sigma}$ (*i.e.*, a is a coatom).

Hence, obviously $\mathcal{L}_1 \circ \mathcal{L}_2$ has the covering property.

($\Rightarrow$): Let $\mathcal{L} \in \mathsf{S}(\mathcal{L}_1, \mathcal{L}_2)$ with the covering property, and $R \subseteq \mathbf{\Sigma}$.

(1): By Lemma 3.2 and Axiom P2, if $R \notin \Xi$ (*i.e.*, there are p^1, $p^2 \in R$ with $p^1_i = p^2_i$ for $i = 1$ or 2), then $\bigvee R \in \mathcal{L}_1 \owedge \mathcal{L}_2$.

(2): Suppose now that $R \in \Xi$. By Lemma 2.9 part 1, if $|R| \leq 2$, then $R \in \mathcal{L}_1 \owedge \mathcal{L}_2$, hence $R \in \mathcal{L}$. Moreover, if $|R| \geq 3$, then for all $s \in \mathbf{\Sigma}$ not in R with $s_1 \in \pi_1(R)$ or $s_2 \in \pi_2(R)$, we have, by Lemma 3.2, $\bigvee(R \bigcup s) = 1$.

(3) **Claim**: Suppose that $R \in \Xi$ and that $|R| = 3$. Write $a := \bigvee R$. Then $a \neq 1$

Proof: Write $R = \{p, q, r\}$ and suppose that $a = 1$. As we have seen $R^0 := \{p, q\} \in \mathcal{L}$. Hence $1 = a = r \vee R^0 \supsetneq p_1 \times \Sigma_2 \cup \Sigma_1 \times q_2 \supsetneq R^0$, a contradiction since $\mathcal{L}$ has the covering property. This proves the claim.

As a consequence, from (2), $1 \gtrdot a$. Moreover, $a \in \Xi$. We write a as $a = \{p_1 \times f(p_1) \mid p_1 \in \pi_1(a)\}$. Hence, f is injective.

(4) **Claim**: $a = R$.

Proof: If $|\Sigma_1| = 3$ or $|\Sigma_2| = 3$, the proof is finished. So we can assume that $|\Sigma_i| \geq 5$ for $i = 1, 2$. Note that any bijection of Σ_2 induces an automorphism of $\mathcal{L}_2$. Suppose that $a \neq R$, hence $|a| \geq 4$. Let $v_2 \in \mathsf{Aut}(\mathcal{L}_2)$ such that its restriction to $\pi_2(a)$ is different from the identity, and with at least three fixed points in $\pi_2(a)$. By Axiom P4, there is $u \in \mathsf{Aut}(\mathcal{L})$ such that on $\mathbf{\Sigma}$, u equals $\mathsf{id} \times v_2$. Hence $c := \{(p_1, v_2 \circ f(p_1)) \mid p_1 \in \pi_1(a)\} \in \mathcal{L}$ and $c \cap a \neq a$, therefore 1 does not cover $c \cap a$. Moreover $|c \cap a| \geq 3$; whence a contradiction by (3). This proves the claim.

(5) Finally, suppose that $R \in \Xi$ and $|R| \geq 4$. Let $b = \bigvee R$. To see that $b = 1$, let $R_0 \in \Xi$ with $R_0 \subseteq R$ and $|R_0| = 3$. By what precedes, $R_0 \in \mathcal{L}$ and $1 \gtrdot R_0$. As a consequence, $b = 1$. □

6. An example with the covering property

Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be complex Hilbert spaces and $T = \mathsf{U}(\mathcal{H}_1) \times \mathsf{U}(\mathcal{H}_2)$. In this section, we give an example in $\mathcal{S}_T(\mathsf{P}(\mathcal{H}_1), \mathsf{P}(\mathcal{H}_2))$, denoted by $\mathsf{P}(\mathcal{H}_1) \textcircled{\Downarrow} \mathsf{P}(\mathcal{H}_2)$, which has the covering property. Moreover, $\mathsf{P}(\mathcal{H}_1) \textcircled{\Downarrow} \mathsf{P}(\mathcal{H}_2)$ is coatomistic, but, as expected, the dual has not the covering property, *i.e.* $\mathsf{P}(\mathcal{H}_1) \textcircled{\Downarrow} \mathsf{P}(\mathcal{H}_2)$ *is not a DAC-lattice.*

Definition 6.1. Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be complex Hilbert spaces, $\mathbf{\Sigma} = \Sigma_{\mathcal{H}_1} \times \Sigma_{\mathcal{H}_2}$, $\mathcal{L}_1 = \mathsf{P}(\mathcal{H}_1)$, $\mathcal{L}_2 = \mathsf{P}(\mathcal{H}_2)$, and $\Sigma_\otimes$ the set of one-dimensional subspaces of $\mathcal{H}_1 \otimes \mathcal{H}_2$. For $V \in \mathsf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$, define $\Sigma_{\Downarrow}[V] := \{(p_1, p_2) \in \mathbf{\Sigma} \mid p_1 \otimes p_2 \in V\}$. Then, $\mathcal{L}_1 \textcircled{\Downarrow} \mathcal{L}_2 := \{\Sigma_{\Downarrow}[V] \mid V \in \mathsf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)\}$, ordered by set-inclusion. For $A \subseteq \mathbf{\Sigma}$, we write $A^\perp := \{q \in \Sigma_\otimes \mid \langle q, p_1 \otimes p_2 \rangle = 0, \forall (p_1, p_2) \in A\}$, where $\langle - | - \rangle$ denotes the scalar product in $\mathcal{H}_1 \otimes \mathcal{H}_2$. Moreover, we denote the set of antilinear maps from $\mathcal{H}_1$ to $\mathcal{H}_2$ by $\mathcal{A}(\mathcal{H}_1, \mathcal{H}_2)$.

Proposition 6.2. *Let m and n be integers, $\mathcal{H}_1 = \mathbb{C}^m$, $\mathcal{H}_2 = \mathbb{C}^n$, $\mathcal{L}_1 = \mathsf{P}(\mathcal{H}_1)$, $\mathcal{L}_2 = \mathsf{P}(\mathcal{H}_2)$, and $\mathbf{\Sigma} = \Sigma_{\mathcal{H}_1} \times \Sigma_{\mathcal{H}_2}$. For $A \in \mathcal{A}(\mathcal{H}_1, \mathcal{H}_2)$, define $X_A \subseteq \mathbf{\Sigma}$ as $X_A := \bigcup \{p_1 \times (A(p_1)^{\perp_2}) \mid p_1 \in \Sigma_1\}$. Then,*

$$\mathcal{L}_1 \textcircled{\Downarrow} \mathcal{L}_2 = \{\bigcap \omega \mid \omega \subseteq \{X_A \mid A \in \mathcal{A}(\mathcal{H}_1, \mathcal{H}_2)\}\}.$$

Proof. Let $\{e_i^1\}_{1 \leq i \leq m}$ and $\{e_j^2\}_{1 \leq j \leq n}$ denote the canonical basis of $\mathbb{C}^m$ and $\mathbb{C}^n$ respectively.

Let $v \in \mathbb{C}^m \otimes \mathbb{C}^n$ and $p = (p_1, p_2) \in \mathbf{\Sigma}$ with $p_1 = \mathbb{C}w_1$ and $p_2 = \mathbb{C}w_2$. Write v, w_1 and w_2 as

$$v = \sum_{i=1}^{m} \sum_{j=1}^{n} s_{ij} e_i^1 \otimes e_j^2, \; w_1 = \sum_{i=1}^{m} \lambda_i e_i^1, \text{ and } w_2 = \sum_{j=1}^{n} \mu_j e_j^2.$$

Let $\lambda = (\lambda_1, \cdots, \lambda_m)^T$ and $\mu = (\mu_1, \cdots, \mu_n)^T$. Let S be the $m \times n$ matrix defined as $S_{ij} = s_{ij}$. Then $p \in \Sigma_{\Downarrow}[v^\perp]$ if and only if $\langle w_1 \otimes w_2, v \rangle = 0$, hence if and only if $\overline{\mu}^T S^T \overline{\lambda} = 0$. Let A be the antilinear map defined by the matrix $S^\perp$. Then, $p \in \Sigma_{\Downarrow}[v^\perp]$ if and only if $p_2 \in A(p_1)^\perp$, that is if and only if $p \in X_A$. As a consequence, $\Sigma_{\Downarrow}[v^\perp] = X_A$.

On the other hand, if $A \in \mathcal{A}(\mathcal{H}_1, \mathcal{H}_2)$, then $X_A = \Sigma_{\Downarrow}[v^\perp]$, where v is given by the formula above with $s_{ij} = (A^T)_{ij}$. □

Remark 6.3. Let $\mathcal{H}$ be a complex Hilbert space of dimension ≥ 3. Then, by Wigner's theorem (see [3], Theorem 14.3.6), any ortho-automorphism of $\mathsf{P}(\mathcal{H})$ is induced by a unitary or antiunitary map on $\mathcal{H}$. Note that if v_1 is a unitary map on $\mathcal{H}_1$ and v_2 is an antiunitary map on $\mathcal{H}_2$, then $v = v_1 \times v_2$ does not induce an automorphism of $\mathcal{L}_1 \textcircled{\Downarrow} \mathcal{L}_2$. Indeed, let X_A be a coatom. Then

$$v(X_A) = \bigcup_{p_1 \in \Sigma_1} v_1(p_1) \times v_2(A(p_1)^{\perp_2}) = \bigcup_{p_1 \in \Sigma_1} p_1 \times ((v_2 \circ A \circ v_1^{-1}(p_1))^{\perp_2}).$$

Now, since A and v_2 are antilinear and v_1 is linear, it follows that $v_2 \circ A \circ v_1^{-1}$ is linear, hence $v(X_A)$ is not a coatom of $\mathcal{L}_1 \textcircled{\Downarrow} \mathcal{L}_2$.

Theorem 6.4. *Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be complex Hilbert spaces, $\mathbf{\Sigma} = \Sigma_{\mathcal{H}_1} \times \Sigma_{\mathcal{H}_2}$, $\mathcal{L}_1 = \mathsf{P}(\mathcal{H}_1)$, $\mathcal{L}_2 = \mathsf{P}(\mathcal{H}_2)$, and $T = \mathsf{U}(\mathcal{H}_1) \times \mathsf{U}(\mathcal{H}_2)$. Then*

(1) *for all $A \subseteq \mathbf{\Sigma}$, we have $\bigvee_{\textcircled{\Downarrow}} A = \Sigma_{\Downarrow}[A^{\perp\perp}]$,*

(2) $\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2 \in \mathcal{S}_T(\mathcal{L}_1, \mathcal{L}_2)$,

(3) $\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2$ *has the covering property and is coatomistic, but, if* $\mathcal{L}_1 \neq 2 \neq \mathcal{L}_2$ *(i.e., the dimension of* $\mathcal{H}_1$ *and* $\mathcal{H}_2$ *is* ≥ 2*), the dual has not the covering property,*

(4) $\mathcal{L}_1 \owedge \mathcal{L}_2 = \{\Sigma_{\Downarrow}[V] \mid V \in \mathsf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2),\, V =\Sigma_{\Downarrow}[V]^{\perp\perp}, V^{\perp} =\Sigma_{\Downarrow}[V^{\perp}]^{\perp\perp}\}$ *(i.e., both* V *and* $V^{\perp}$ *are spanned by product vectors),*

(5) *if* $\mathcal{L}_1 \neq 2 \neq \mathcal{L}_2$*, then* $\mathcal{L}_1 \owedge \mathcal{L}_2 \subsetneqq \mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2 \subsetneqq \mathcal{L}_1 \ovee \mathcal{L}_2$*.*

Proof. (1): This follows directly from Definition 6.1.

(2): Obviously, $\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2$ is a simple closure space on $\mathbf{\Sigma}$. Let $a \in \mathcal{L}_1 \owedge \mathcal{L}_2$. By Definition, $a^{\#} =\Sigma_{\Downarrow}[a^{\perp}]$. Hence, $a^{\#} \subseteq a^{\perp}$, $a^{\perp\perp} \subseteq a^{\#\perp}$, thus $\Sigma_{\Downarrow}[a^{\perp\perp}] \subseteq a^{\#\#} = a$, and therefore $a =\Sigma_{\Downarrow}[a^{\perp\perp}]$. As a consequence, $\mathcal{L}_1 \owedge \mathcal{L}_2 \subseteq \mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2$.

Let $V \in \mathsf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$ such that $\Sigma_{\Downarrow}[V] = p_1 \times A_2$. Then

$$p_1 \otimes (A_2^{\perp_2\perp_2}) = (p_1 \otimes A_2)^{\perp\perp} \subseteq V,$$

therefore $A_2^{\perp_2\perp_2} \subseteq A_2$, hence $A_2 \in \mathcal{L}_2$. As a consequence, Axiom P3 holds.

Axiom P4 with $T = \mathsf{U}(\mathcal{H}_1) \times \mathsf{U}(\mathcal{H}_2)$ holds in $\mathsf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$, therefore obviously also in $\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2$.

(3): The covering property holds in $\mathsf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$ (see [6], Theorem 34.2), hence, by (1), also in $\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2$. Moreover, since $\mathsf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$ is coatomistic, so is $\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2$.

Next, let $p \in \mathbf{\Sigma}$. Then $x =\Sigma_{\Downarrow}[p^{\perp}] = p^{\#}$ is a coatom of $\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2$. Now, there is $R \in \Xi(\mathbf{\Sigma})$ (see Definition 3.5) with $|R| = 2$, such that $x \cap R = \emptyset$. By Lemma 2.9, $R \in \mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2$. Moreover $x \bigvee_{\textcircled{$\Downarrow$}} R = 1$ since x is a coatom. Hence, writing $R = \{p, q\}$, and the order relation, meet, join, bottom and top elements in $(\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2)^*$ by $\leq_*$, $\bigwedge_*$, $\bigvee_*$, 0_*, and 1_* respectively, we find that $x \bigwedge_* R = 0_*$ and $x \bigvee_* R = 1_* \gneqq_* p \gneqq_* R$. Therefore, $(\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2)^*$ does not have the covering property.

(4) Let $p \in \mathbf{\Sigma}$ and $q \in p^{\#\perp}$. Write $q = \mathbb{C}v$ with $v \in \mathcal{H}_1 \otimes \mathcal{H}_2$. For $i = 1$ and $i = 2$, let $\{w_i^k\}$ be an ortho-basis of $p_i^{\perp_i}$, and let $x_i \in p_i$ (*i.e.*, $p_i = \mathbb{C}x_i$). Then v can be decomposed as

$$v = \alpha x_1 \otimes x_2 + \sum_{k_2} \beta_{k_2} x_1 \otimes w_2^{k_2} + \sum_{k_1} \beta_{k_1} w_1^{k_1} \otimes x_2 + \sum_{l_1 l_2} \gamma_{l_1 l_2} w_1^{l_1} \otimes w_2^{l_2}.$$

Now, $p^{\#} = p_1^{\perp_1} \times \Sigma_{\mathcal{H}_2} \cup \Sigma_{\mathcal{H}_1} \times p_2^{\perp_2}$. Hence, since $q \in p^{\#\perp}$, we find that $\gamma_{l_1 l_2} = \beta_{k_1} = \beta_{k_2} = 0$, for all k_1, k_2, l_1 and l_2. Therefore, $v \in p$, hence $p^{\#\perp} = p$.

Let $a \in \mathcal{L}_1 \owedge \mathcal{L}_2$. From (1), $a =\Sigma_{\Downarrow}[a^{\perp\perp}]$. On the other hand, $a^{\#} =\Sigma_{\Downarrow}[a^{\perp}]$. Now, $a = a^{\#\#} = \bigcap\{p^{\#} \mid p \in a^{\#}\}$. Hence, by the preceeding,

$$a^{\perp} = \left(\bigcup\{p^{\#\perp} \mid p \in a^{\#}\}\right)^{\perp\perp} = a^{\#\perp\perp},$$

therefore $a^{\perp}$ is also spanned by product vectors. Hence, writing $V = a^{\perp\perp}$, we find that $a =\Sigma_{\Downarrow}[V]$, $V =\Sigma_{\Downarrow}[V]^{\perp\perp}$ and $V^{\perp} =\Sigma_{\Downarrow}[V^{\perp}]^{\perp\perp}$.

Let $V \in \mathsf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$ such that both V and $V^{\perp}$ are spanned by product vectors. Let $a :=\Sigma_{\Downarrow}[V]$ and $b :=\Sigma_{\Downarrow}[V^{\perp}]$. Since $\Sigma_{\Downarrow}[V]^{\perp\perp} = V$ and also $\Sigma_{\Downarrow}[V^{\perp}]^{\perp\perp} = V^{\perp}$, then $\Sigma_{\Downarrow}[\Sigma_{\Downarrow}[V]^{\perp}] =\Sigma_{\Downarrow}[V^{\perp}]$ and $\Sigma_{\Downarrow}[\Sigma_{\Downarrow}[V^{\perp}]^{\perp}] =\Sigma_{\Downarrow}[V]$. Therefore $a^{\#} = b$ and $b^{\#} = a$. As a consequence, $a \in \mathcal{L}_1 \owedge \mathcal{L}_2$.

(5): By (4), $\mathcal{L}_1 \owedge \mathcal{L}_2 \neq \mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2$. On the other hand, by Theorem 5.4, $\mathcal{L}_1 \ovee \mathcal{L}_2$ does not have the covering property, whereas by (3), $\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2$ has the covering property. As a consequence, $\mathcal{L}_1 \mathbin{\textcircled{$\Downarrow$}} \mathcal{L}_2 \neq \mathcal{L}_1 \ovee \mathcal{L}_2$. □

REFERENCES

[1] Aerts, D.: Description of many separated physical entities without the paradoxes encountered in quantum mechanics. Found. Phys. **12**, 1131–1170 (1982)

[2] Aerts, D.: The missing elements of reality in the description of quantum mechanics of the EPR paradox situation. Helv. Phys. Acta **57**, 421–428 (1984)

[3] Faure, C.A., Frölicher, A.: Modern projective geometry. Mathematics and its Applications, vol. 521. Kluwer Academic Publishers, Dordrecht (2000)

[4] Ischi, B.: Orthocomplementation and compound systems. Internat. J. Theoret. Phys. **44**, 2207–2217 (2005)

[5] Ischi, B.: Weak tensor products of complete atomistic lattices. Algebra Universalis **57**, 239–258 (2007)

[6] Maeda, F., Maeda, S.: Theory of symmetric lattices. Die Grundlehren der mathematischen Wissenschaften, vol. 173. Springer-Verlag, New York (1970)

[7] Piron, C.: Axiomatique quantique. Helv. Phys. Acta **37**, 439–468 (1964) (French)

[8] Piron, C.: Foundations of quantum physics. Mathematical Physics Monograph Series, vol. 19. W. A. Benjamin, Inc., Reading, Mass.-London-Amsterdam (1976)

[9] Pulmannová, S.: Tensor product of quantum logics. J. Math. Phys. **26**, 1–5 (1985)

E-mail address: `boris.ischi@edu.ge.ch`

COLLÈGE DE CANDOLLE, 5 RUE D'ITALIE, 1204 GENEVA, SWITZERLAND